

Molecular Gas in nearby Early-Type Powerful Classical Radio Galaxies

Stéphane Leon

Instituto de Astrofísica de Andalucía, Camino Bajo de Huétor, 24 Apdo 3004, 18080 Granada, Spain

Jeremy Lim

Institute of Astronomy & Astrophysics, Academia Sinica, P.O. Box 23-141, Taipei 106, Taiwan

Françoise combes

Observatoire de Paris, LERMA, 61 Avenue de l'Observatoire, 75014 Paris, France

Dihn-V-Trung

Institute of Astronomy & Astrophysics, Academia Sinica, P.O. Box 23-141, Taipei 106, Taiwan

Abstract. We report a survey for molecular gas in nearby powerful radio galaxies. Eight of the eighteen radio galaxies observed were detected with molecular masses in the range 10^7 – 10^9 $M_{\odot}$, similar to the same survey we performed towards 3C radio galaxies. The upper limits of molecular gas in the remainder are typically of 10^8 $M_{\odot}$, indicating that very few radiogalaxies have molecular gas reservoir with more than 10^9 $M_{\odot}$.

1. Introduction

Powerful classical double-lobed radio galaxies are (at low redshifts) hosted almost exclusively by luminous early-type galaxies. These galaxies are usually elliptical galaxies. Given that normal elliptical galaxies usually exhibit undetectable quantities of cool (neutral) gas, what fuels the central supermassive black holes in elliptical radio galaxies? Our ^{12}CO survey of a complete sample of 3C galaxies at low redshift (Lim et al., this volume) reveals that the vast majority have 5σ upper limits in molecular gas masses of $2\text{--}5 \times 10^8$ $M_{\odot}$ (for a linewidth of 500 km/s and a Galactic CO-to- H_2 conversion factor). To better understand the molecular gas content of classical radio galaxies, we are undertaking a survey of nearby early-type powerful radio galaxies. These galaxies were originally selected for study by Verdoes Kleijn et al. (1999) with the HST, and were drawn from a catalog of one-hundred seventy-six radio-loud galaxies constructed by Condon & Broderick (1988).

2. Results

Of the eighteen galaxies where sensitive 12CO measurements have now been made, we detected eight (i.e., a detection rate of 45%). The galaxies have molecular gas masses in the range $10^7 - 10^9 M_\odot$. All the galaxies detected have double-horned 12CO line profiles characteristic of a rotating disk or torus, commensurate with the disk-like dust features seen in these galaxies. In the case of 3C 31, our follow-up imaging observations with the IRAM PdBI confirm that the molecular gas is distributed in a rotating disk/torus with size similar to its dust disk. These results, summarized in Table 1, reinforce our argument (Lim et al. 2000) that, in the vast majority of cases, the molecular gas in powerful classical radio galaxies most likely originates from a merger between a pre-existing elliptical galaxy and a relatively less massive gas-rich (disk) galaxy. Radio activity is triggered only when sufficient gas reaches the nucleus of the elliptical galaxy, thereby explaining the relatively relaxed nature of both the accreted molecular gas and parent elliptical galaxy.

Table 1. Results of CO survey of nearby powerful radio galaxies

Name	Redshift (z)	IRAS	Optical Dust	$M(\text{H}_2)$ ($M_\odot$)
NGC 193	0.014		✓	$< 1.2 \times 10^8$
NGC 315	0.016	✓	✓	$(3.03 \pm 0.34) \times 10^8$
NGC 383 (3C 31)	0.017	✓	✓	$(1.06 \pm 0.04) \times 10^9$
NGC 541	0.018		✓	$(2.47 \pm 0.38) \times 10^8$
UGC 1841 (3C 66B)	0.021			$< 2.8 \times 10^8$
NGC 2329	0.019	✓		$< 1.4 \times 10^8$
NGC 2892	0.023			$< 3.4 \times 10^8$
NGC 3801	0.011		✓	$(3.77 \pm 0.22) \times 10^8$
NGC 3862 (3C 264)	0.022	✓	✓	$(3.09 \pm 0.26) \times 10^8$
NGC 4261 (3C 270)	0.007	✓	✓	$< 3.6 \times 10^7$
NGC 4335	0.015		✓	$< 1.2 \times 10^8$
NGC 4374 (3C 272.1)	0.003	✓	✓	$(1.40 \pm 0.38) \times 10^7$
NGC 4486 (3C 274)	0.007	✓		standing wave ripples
NGC 5127	0.016		✓	$< 1.4 \times 10^8$
NGC 5141	0.018		✓	$< 4.4 \times 10^8$
NGC 7052	0.016	✓		$(3.45 \pm 0.23) \times 10^8$
UGC 12064 (3C 449)	0.018	tentative	✓	$(2.85 \pm 0.23) \times 10^8$
NGC 7626	0.011		✓	$< 0.8 \times 10^8$

References

- Condon, J. J.; Broderick, J. J. 1988, AJ, 96, 30
 Lim, J., Leon, S., Combes, F., Dinh-V-Trung 2000, ApJ, 545, L93
 Verdoes Kleijn, G. A., Baum, S. A., de Zeeuw, P.T., O'Dea, C. 1999, AJ, 118, 2592